

AOFAS Lesser Metatarsophalangeal-Interphalangeal Scale

Patient Name: _____

Patient MRN: _____

Date: _____

I. Pain (40 points)

<input type="checkbox"/> None	+40
<input type="checkbox"/> Mild, occasional	+30
<input type="checkbox"/> Moderate, daily	+20
<input type="checkbox"/> Severe, almost always present	+0

II. Function (45 points)

Activity limitations

<input type="checkbox"/> No limitations	+10
<input type="checkbox"/> No limitation of daily activities, limitation of recreational activities	+7
<input type="checkbox"/> Limited daily and recreational activities	+4
<input type="checkbox"/> Severe limitation of daily and recreational activities	+0

Footwear requirements

<input type="checkbox"/> Fashionable, conventional shoes, no insert required	+10
<input type="checkbox"/> Comfort footwear, shoe insert	+5
<input type="checkbox"/> Modified shoes or brace	+0

MTP joint motion (dorsiflexion plus plantarflexion)

<input type="checkbox"/> Normal or mild restriction (75° or more)	+10
<input type="checkbox"/> Moderate restriction (30° - 74°)	+5
<input type="checkbox"/> Severe restriction (less than 30°)	+0

IP joint motion (plantarflexion)

<input type="checkbox"/> No restriction	+5
<input type="checkbox"/> Severe restriction (less than 10°)	+0

MTP-IP stability (all directions)

<input type="checkbox"/> Stable	+5
<input type="checkbox"/> Definitely unstable or able to dislocate	+0

Callus related to lesser MTP-IP

<input type="checkbox"/> No callus or asymptomatic callus	+5
<input type="checkbox"/> Callus, symptomatic	+0

III. Alignment (15 points)

<input type="checkbox"/> Good, lesser toes well aligned	+15
<input type="checkbox"/> Fair, some degree of lesser toe malalignment observed, no symptoms	+8
<input type="checkbox"/> Poor, severe malalignment, symptoms	+0

IV. Total Score (100 points):

_____ Pain Points +

_____ Function Points +

_____ Alignment Points =

_____ Total Points/100 points